

# Coronavirus COVID-19

BC Centre for Disease Control | BC Ministry of Health



## Infection Prevention and Control (IPC) Protocol for Obstetrical Procedures During COVID-19

May 24, 2020

### Guiding Principles:

- Provider Safety
- Patient Safety
- PPE Conservation

### Approach to IPC Includes:

- Patient COVID-19 Assessment
- Surgical Risk Assessment
- PPE Recommendation
- PPE Allocation Framework<sup>1</sup>

## Background/Current Status

This guidance supports B.C. health authorities with ongoing obstetrical operative procedures in the context of the COVID-19 pandemic.

Through effective public health measures the COVID-19 pandemic curve has reached its peak and is on the downward slope. The prevalence of COVID within the maternity population in BC is low and in keeping with the general population. The current recommendations are based on the low prevalence within the pregnancy population and our current understanding of risk.

**Based on the epidemiology of COVID-19 in B.C.<sup>2</sup>, obstetrical patients who do not have risk factors or symptoms of COVID-19 should not be considered suspect cases.** This is based on the advice of the BC Centre for Disease Control (BCCDC), the Office of the Provincial Health Officer (PHO) and the Provincial Infection Control Network of BC (PICNet). BCCDC, PHO, maternity consultants and PICNet review the epidemiology on a regular basis and will amend or update this advice as required.

**All patients and support persons arriving to a maternity or birthing unit must be assessed for risk factors and symptoms of COVID-19, and where appropriate tested.** Refer to the most recent BCCDC testing and

<sup>1</sup> COVID-19: Emergency Prioritization in a Pandemic Personal Protective Equipment (PPE) Allocation Framework, Provincial COVID-19 Task Force, March 25, 2020: <https://www2.gov.bc.ca/assets/gov/health/about-bc-s-health-care-system/office-of-the-provincial-health-officer/covid-19/ppe-allocation-framework-march-25-2020.pdf>

<sup>2</sup> Epidemiologic considerations: daily case counts; test positivity rate; incidence rate; point prevalence.



Ministry of Health



BC Centre for Disease Control

If you have fever, a new cough, or are having difficulty breathing, call 8-1-1.



laboratory guidance for COVID-19. For women who are confirmed cases of COVID-19 please see the latest BCCDC recommendations for guidance on self-isolation and management during pregnancy. PCR, NAT, or serologic SARS-CoV-2 tests are not currently recommended for screening asymptomatic patients due to the low clinical sensitivity and low positive predictive value in lower acuity patients and in low prevalence contexts.

The guidance includes a patient screening tool and classification of patients based on a Patient Risk Categorization into green, yellow and red categories. **The entire surgical team, including anesthesiologist, surgeon, assistant, nurses, etc., is responsible for deciding the Patient Risk Category together** and the expectation is that obstetrical surgery will not be delayed as a result.

Given this guidance and the current low incidence and prevalence of COVID-19 in B.C, the risk of infection or transmission to health care workers when protocols are followed is extremely low.

## Scope

---

This guidance and protocol does not apply to non-obstetrical populations. There is separate provincial guidance available regarding specific adult and pediatric surgical protocols.

### A. Pre-surgical Patient Assessment

---

- For patients presenting for scheduled obstetric surgery, the **COVID-19 Surgical Patient Assessment Form (see Appendix 1)** should be completed 24 to 72 hours prior to scheduled surgery, by the pre-admission unit (nurse, medical office assistant or anesthesiologist) over the phone, and then repeated in person when the patient arrives at the hospital on the day of surgery. This is done to allow for testing (if clinically indicated) to be complete prior to scheduled procedures<sup>3,4</sup>.
- Procedures performed under local or regional anesthesia should be performed under droplet precautions. For cases where a patient is classified as yellow or red, the risk of conversion to general anesthesia must be discussed at the huddle to help guide appropriate PPE under section D.
- There needs to be a mechanism in place within each facility unit to ensure the Surgical Patient Assessment Form is included in the patient chart.
- For urgent or emergent procedures, the COVID-19 Patient Assessment Form shall be completed upon arrival to the peri-operative area. At many sites, an equivalent site-specific COVID patient assessment form will have been completed on admission to the hospital for labouring patients. This assessment should be considered equivalent to the pre-surgical form due to the evolving nature of obstetrical care. A delay in the OR procedure should not occur if the pre-surgical form (or equivalent) has not been performed due to the time-sensitive nature of obstetrical care. If a patient has signs and symptoms consistent with COVID-19 testing should be conducted as per provincial COVID-19 testing guidelines<sup>3</sup>.
- IPC risk categories have been developed to guide PPE use before, during, and after a surgical procedure:
  - i. Low or no risk (green) – a patient with no risk factors for COVID-19, and/or no symptoms or signs of COVID-19, and/or a negative COVID-19 RNA test where relevant

---

<sup>3</sup> As defined by the BCCDC. See <http://www.bccdc.ca/health-professionals/clinical-resources/covid-19-care/lab-testing> for more information.

<sup>4</sup> Every attempt should be made to assess the patient in their preferred language.

- ii. Unknown risk (yellow) – a patient where the risk factors history and symptomatology are unknown, and a RNA COVID-19 test result is pending or unknown, OR a CODE OB/CODE PINK patients where a history cannot be obtained
- iii. Moderate to high risk (red) – a patient with risk factors for COVID-19, and/or symptoms or signs of COVID-19, and/or a COVID-19 RNA test result is pending or unknown, OR a lab confirmed COVID-19 RNA test.
- For the CODE OB/CODE PINK patient where an appropriate history has been obtained and has not changed, for example, fever in labour, during the admission, patients can continue to be classified as either green or red. **For CODE OB/CODE PINK patients where a history cannot be obtained due to the emergent nature of care they are classified as yellow.**
- Obstetrical cases should not be delayed while COVID-19 test results are pending.

## B. Pre-surgical Procedure Huddle

---

- The pre-surgical huddle, when the full surgical team is engaged (anesthetist, surgeon, maternity care provider, pediatricians, assistants, nurses, etc.), is one of the strongest determinants for achieving the highest levels of safety and quality. All of the other usual elements of a surgical checklist should also be discussed at this time.
- The Patient Risk Category is determined based on information gathered from the assessment form, including COVID-19 testing results, if applicable (see Appendix 1).
- For operative procedures that occur during the course of labour or post partum care, the team needs to ask “**has the patient’s clinical status changed to warrant a change in patient risk category?**” such as new onset of fever in labour (green to red or COVID swab turns negative red to green).
- **Surgical team members must agree on the Patient Risk Category (green, yellow or red)**
- Neuraxial anesthesia is the usual practice for the majority of obstetrical surgical cases. Procedures performed under local or regional anesthesia should be performed under droplet precautions. For cases where a patient is classified as yellow or red, the risk of conversion to general anesthesia must be discussed at the huddle to help guide appropriate use of PPE.
- Recommended PPE to be used during the surgical procedure is provided in **Section D: Algorithm for Management of Obstetrical Surgical Patients.**

## C. Air Clearance

---

- Airflow considerations, including appropriate times for air clearance post-aerosol generating medical procedure (AGMP), should be made for each OR suite in consultation with local infection prevention and control (IPAC), and facilities, maintenance and operations (FMO).
  - In most ORs and post-operative areas, the relative humidity (RH) is kept between 40% and 45% which aids in reducing the amount of virus or bacteria in the air.
  - Raising the RH not only causes more rapid “fallout” of particles below the respiratory zone, but also has been documented to be beneficial for clearing respiratory secretions and hydrating mucous membranes with associated improved outcome.
  - Increased RH decreases viral survival. The air exchange rate (or air changes per hour – ACH) is kept between 18 and 23 in most ORs (higher in positive pressure rooms).
  - Between the increased RH and the ACH, the potential for bioaerosol spread will be reduced by over 95% within 10-12 minutes following aerosol creation (extubation).
- The AGMP should be performed with the door(s) closed. Limiting the number of personnel and equipment in the room and minimizing door openings is a key element in environmental infection control.

## E. Algorithm for Management of Obstetrical Surgical Patients

Infection Prevention & Control Risk Category			
	Green	Yellow <i>(Code OB or Code Pink)</i>	Red
<b>Team Huddle</b>	Team to Review: <ul style="list-style-type: none"> <li>Confirm patient Risk Category</li> <li>Anesthetic approach</li> <li>Staff to be in OR (eg. RM/FP)</li> <li>Presence of support person in OR as per routine</li> </ul>	Team to Review: <ul style="list-style-type: none"> <li>Confirm patient Risk Category</li> <li>Anesthetic approach</li> <li>Staff to be in OR (eg RM/FP)</li> <li>Presence of support person in OR*</li> </ul>	Team to Review: <ul style="list-style-type: none"> <li>Confirm patient Risk Category</li> <li>Anesthetic approach</li> <li>Staff to be in OR (eg RM/FP)</li> <li>Presence of support person in OR*</li> </ul>
<b>Neuraxial Anesthesia</b>	Routine personnel in the OR <ul style="list-style-type: none"> <li>All personnel in the OR:                             <ul style="list-style-type: none"> <li>Routine OR PPE</li> </ul> </li> </ul>	Limit personnel in the OR <ul style="list-style-type: none"> <li>Surgical Team and Anesthesia:                             <ul style="list-style-type: none"> <li>Fit-tested N95 Respirator</li> <li>Face shield or goggles</li> <li>Gown &amp; gloves</li> </ul> </li> </ul> Pediatric Team and other personnel in droplet and contact in room at start of procedure  If appropriate neuraxial analgesia can consider droplet and contact for all personnel	Limit personnel in the OR <ul style="list-style-type: none"> <li>Surgical Team and Anesthesia:                             <ul style="list-style-type: none"> <li>Fit-tested N95 Respirator</li> <li>Face shield or goggles</li> <li>Gown &amp; gloves</li> </ul> </li> </ul> Pediatric Team and other personnel in droplet and contact in room at start of procedure
<b>IF General Anesthetic: Intubation and Extubation</b>	Routine personnel in the OR <ul style="list-style-type: none"> <li>All personnel in the OR:                             <ul style="list-style-type: none"> <li>Routine OR PPE</li> </ul> </li> </ul>	Limit personnel in the OR <ul style="list-style-type: none"> <li>All staff in the OR don:                             <ul style="list-style-type: none"> <li>Fit-tested N95 Respirator</li> <li>Face shield or goggles</li> <li>Gown &amp; gloves</li> </ul> </li> </ul> Pediatric Team in N95 in room at start of procedure  All non-essential personnel to leave the room for extubation	Limit personnel in the OR <ul style="list-style-type: none"> <li>All staff in the OR:                             <ul style="list-style-type: none"> <li>Fit-tested N95 Respirator</li> <li>Face shield or goggles</li> <li>Gown &amp; gloves</li> </ul> </li> </ul> Pediatric Team in N95 in room at start of procedure  All non-essential personnel to leave the room for extubation
<b>Recovery Regional</b>	Recover as per routine at site	<ul style="list-style-type: none"> <li>Recover in the designated COVID location using Droplet/Contact Precautions until ready to move to designated unit</li> </ul>	<ul style="list-style-type: none"> <li>Recover in the designated COVID location using Droplet/Contact Precautions until ready to move to designated unit</li> </ul>
<b>Recovery GA</b>	Recover as per routine at site	<ul style="list-style-type: none"> <li>Recover in the OR suite until ready to move to designated unit</li> </ul>	<ul style="list-style-type: none"> <li>Recover in the OR suite until ready to move to designated unit</li> </ul>
<b>Cleaning and Disinfection</b>	Cleaning should be determined as per site specific routine protocols	All cleaning staff in OR don: <ul style="list-style-type: none"> <li>Surgical mask</li> <li>Eye protection</li> <li>Gown/Gloves</li> </ul>	All cleaning staff in OR don: <ul style="list-style-type: none"> <li>Surgical mask</li> <li>Eye protection</li> <li>Gown/Gloves</li> </ul>
<b>Disposition</b>	Transfer patient to postpartum as per routine care.	Return patient to appropriate inpatient unit based on further patient risk assessment.	Return to appropriate COVID-19 isolation room if confirmed positive or isolation room if unknown.

\* The guiding principle is that there is a support person in the OR if the procedure is performed under neuraxial anesthesia, including RED and YELLOW patients. Support persons should be screened for symptoms and wear appropriate PPE as per site/HA protocol.

## Appendix 1: COVID-19 Surgical Patient Assessment Form - Obstetrics

Health Authority LOGO

### Patient Information

Name:  
Date of Birth:  
Language:  
PHN:

#### NURSE OR MEDICAL OFFICE ASSISTANT SCREEN:

Able to obtain patient history?  Yes  No If No, go to Physician Screen section

#### ***Does the patient have a risk factor for COVID-19 exposure? In the last 14 days has the patient:***

Returned from travel outside of Canada?  Yes  No When? Date: \_\_\_\_\_

Been in close contact with anyone diagnosed with lab confirmed COVID-19?  Yes  No When? Date: \_\_\_\_\_

Lived or worked in a setting that is part of a COVID-19 outbreak?  Yes  No When? Date: \_\_\_\_\_

Been advised to self-isolate or quarantine at home by public health?  Yes  No Contact info: \_\_\_\_\_

#### ***Does the patient have new onset COVID-19 like symptoms in the last 14 days?***

24 to 72 hours prior – Date/Time: \_\_\_\_\_ Day of surgery – Date/Time: \_\_\_\_\_

Fever  Yes  No Fever  Yes  No

Cough  Yes  No Cough  Yes  No

Shortness of breath  Yes  No Shortness of breath  Yes  No

Diarrhea  Yes  No Diarrhea  Yes  No

Nausea and/or vomiting  Yes  No Nausea and/or vomiting  Yes  No

Headache  Yes  No Headache  Yes  No

Runny nose/nasal congestion  Yes  No Runny nose/nasal congestion  Yes  No

Sore throat or painful swallowing  Yes  No Sore throat or painful swallowing  Yes  No

Loss of sense of smell  Yes  No Loss of sense of smell  Yes  No

Loss of appetite  Yes  No Loss of appetite  Yes  No

Chills  Yes  No Chills  Yes  No

Muscle aches  Yes  No Muscle aches  Yes  No

Fatigue  Yes  No Fatigue  Yes  No

Screened by:

Signature:

Screened by:

Signature:

**PHYSICIAN/SURGEON SCREEN:**

COVID-19 NP test performed  Yes  No Date: \_\_\_\_\_  
 Result:  Negative  Positive

If test has not been performed, do you recommend testing patient?  Yes  No Reason: \_\_\_\_\_

Unable to perform swab?  Yes  No Reason: \_\_\_\_\_

**Screened by:** \_\_\_\_\_ **Signature:** \_\_\_\_\_ **Date/Time:** \_\_\_\_\_

**FINAL SURGICAL TEAM ASSESSMENT:**

COVID-19 risk factor (travel, contact, outbreak)?  Yes  No  Unknown

COVID-19 like symptoms that cannot be explained by another medical or surgical diagnosis?  Yes  No  Unknown

COVID-19 test result?  Yes  No  Unknown  N/A

**PATIENT RISK CATEGORY TABLE:**


COVID-19 Risk Factors	COVID-19 Symptoms	COVID -19 Test Results	COVID-19 Risk Category
NO	NO	NOT REQUIRED	GREEN
NO	NO	NEGATIVE	GREEN
YES	NO	NEGATIVE	GREEN
NO	UNKNOWN	NEGATIVE	GREEN
NO	YES	NEGATIVE	GREEN
YES	YES	NEGATIVE	GREEN
UNKNOWN	UNKNOWN	UNKNOWN/PENDING	YELLOW
YES	NO	UNKNOWN/PENDING	RED
NO	YES	UNKNOWN/PENDING	RED
YES	YES	UNKNOWN/PENDING	RED
-	-	<b>POSITIVE</b>	RED

**PATIENT RISK CATEGORY (CIRCLE ONE):**

GREEN	YELLOW	RED
-------	--------	-----

**Key Informants**

---



# Coronavirus COVID-19

BC Centre for Disease Control | BC Ministry of Health



Dr. Chelsea Elwood, BCCW, PHSA (Co-Lead)  
Dr. Ellen Giesbrecht, BCCW PSBC, PHSA (Co-Lead)  
Dr. Titus Wong, PICNet, PHSA (Co-Lead)  
Dr. Michael Murray, CPSBC (Co-Lead)

Dr. Deborah Money, UBC  
Dr. Roanne Preston, UBC, VGH, PHSA  
Tara Donovan, PICnet, PHSA  
Dr. Trevor Corneil, MOH

Members of the Provincial IPC Protocol for Surgical Procedures  
Members of the Perinatal Services BC Steering Committee

To provide feedback on this document please email [covidguidelines@bcpsqc.ca](mailto:covidguidelines@bcpsqc.ca) - including the document title in the subject line.

## References

COVID-19: Emergency Prioritization in a Pandemic Personal Protective Equipment (PPE) Allocation Framework, Provincial COVID-19 Task Force, March 25, 2020  
[https://www2.gov.bc.ca/assets/gov/health/about-bc-s-health-care-system/office-of-the-provincial-health-officer/covid-19/ppe\\_allocation\\_framework\\_march\\_25\\_2020.pdf](https://www2.gov.bc.ca/assets/gov/health/about-bc-s-health-care-system/office-of-the-provincial-health-officer/covid-19/ppe_allocation_framework_march_25_2020.pdf)

Wang W, Xu Y, Gao R, et al. Detection of SARS-CoV-2 in Different Types of Clinical Specimens. JAMA  
Published Online First: Mar. 11, 2020. doi:10.1001/jama.2020.3786

Yang Y, Yang M et al. Evaluating the accuracy of different respiratory specimens in the laboratory diagnosis and monitoring the viral shedding of 2019-nCoV infections MedRxiv Published Online First: Feb. 17, 2020.  
<https://doi.org/10.1101/2020.02.11.20021493>

Jiang G et al. Application and optimization of RT-PCR in diagnosis of SARS-CoV-2 infection  
MedRxiv Feb. 28, 2020. <https://doi.org/10.1101/2020.02.25.20027755>

Ali, J et al. Optimizing diagnostic strategy for novel coronavirus pneumonia, a multi-center study in Eastern China MedRxiv Feb. 17, 2020. <https://doi.org/10.1101/2020.02.13.20022673> Fang Y, Zhang H, Xie J, et al. Sensitivity of Chest CT for COVID-19: Comparison to RT-PCR. Radiology 2020: 200432. doi:10.1148/radiol.2020200432



Ministry of Health



BC Centre for Disease Control

**If you have fever, a new cough, or are having difficulty breathing, call 8-1-1.**



Ng, K., Poon, B. H., Puar, T. H. K., Quah, J. L. S., Loh, W. J., Wong, Y. J., ... Raghuram, J. (2020). COVID-19 and the risk to health care workers: a case report. *Annals of Internal Medicine*.

Yao, W., Wang, T., Jiang, B., Gao, F., Wang, L., Zheng, H., ... Mei, W. (2020). Emergency tracheal intubation in 202 patients with COVID-19 in Wuhan, China: lessons learnt and international expert recommendations. *British Journal of Anaesthesia*.

Cook TM, El-Boghdadly K, McGuire B, McNarry AF, Patel A, Higgs A. Consensus guidelines for managing the airway in patients with COVID-19: Guidelines from the Difficult Airway Society, the Association of Anaesthetists the Intensive Care Society, the Faculty of Intensive Care Medicine and the Royal College of Anaesthetists. *Anaesthesia*. 2020.