



Coronavirus COVID-19

BC Centre for Disease Control | BC Ministry of Health



Clinical Guidance on COVID-19 Vaccines for people with hematological malignancy at any stage of treatment and/or who have undergone hematopoietic stem cell transplant or CAR-T cell therapy in the past 6 months

This guidance is intended for healthcare providers and is based on known evidence as of March 1, 2022.

This document relates to people with hematological malignancies and those who have undergone hematopoietic stem cell transplant or CAR-T cell therapy. Please refer to other guidance document for people with solid cancers. For general information, please refer to BCCDC Guidance and information on [COVID-19 vaccines for providers](#).

Background and Context

Hematologic malignancies (blood cancers such as leukemias, myelodysplastic syndromes, myeloproliferative neoplasms, lymphomas and multiple myeloma) can be treated with chemotherapy, hematopoietic stem cell transplant (HSCT) or chimeric antigen receptor T-cell (CAR-T) therapies with either a curative intent or to prolong survival. The hematologic malignancy itself or the anti-cancer therapies can result in long-lasting immunodeficiency, and COVID-19 infection in this population is associated with a significantly higher risk of hospitalization and death.¹

This guidance is based on a review of the vaccines approved by Health Canada for the prevention of COVID-19 disease caused by the SARS-CoV-2 virus:

- **mRNA vaccines:** tozinameran (COMIRNATY, Pfizer-BioNTech),² elasomeran (SPIKEVAX, Moderna)³
- **Replication-defective adenoviral vector vaccine:** ChADOx1-S (VAXZEVRIA, AstraZeneca),⁴ Ad26.COV2.S (Janssen COVID-19 Vaccine, Janssen)⁵

Currently, anyone in British Columbia who is 5 years and older is eligible for COVID-19 immunization. Health Canada has authorized a COMIRNATY (Pfizer-BioNTech) vaccine made for children aged 5 to 11. NACI has released their statement for this age group⁶.

People who receive the mRNA vaccine (COMIRNATY [Pfizer-BioNTech] or SPIKEVAX [Moderna]) for their first dose, will be offered either mRNA vaccine for subsequent doses, with the exception of preferential recommendations based on age and immunosuppression.^{7,8} BC has taken the proactive step to expand booster doses for all individuals 12 years and older, not just those at high risk. However, it is particularly recommended for individuals 12-17 years of age who are at higher risk of severe illness due to COVID-19.⁹ All booster doses will be mRNA vaccines.¹⁰



Third doses:

To date, people who are moderately to severely immunocompromised have been observed to have generally lower antibody responses and lower vaccine effectiveness from COVID-19 vaccines compared to the general population. The National Advisory Committee on Immunization¹¹ has reviewed this evidence and recent studies that demonstrate that some people who are immunocompromised develop an improved antibody response after a third dose of vaccine.

As such, as of February 3, 2022, people (5 years and older) who are severely immunocompromised in B.C. are eligible to receive a third dose of an mRNA COVID-19 vaccine.¹²

A minimum interval of 28 days between dose 2 and dose 3 is recommended for those eligible for a third dose. As per the BC Immunization Manual, COMIRNATY (Pfizer-BioNTech) is recommended for those 5-11 years of age. For individuals 12 years of age and older, SPIKEVAX (Moderna) is preferred for the third dose; however, if SPIKEVAX (Moderna) is unavailable (or if the individual prefers), the COMIRNATY (Pfizer-BioNTech) vaccine may be provided.^{12,13}

Specifics on current eligibility for a third dose may be reviewed here: <https://www2.gov.bc.ca/gov/content/covid-19/vaccine/register#immunocompromised>

Other vaccines:

The VAXZEVRIA (AstraZeneca) vaccine program has been stopped in B.C. for first doses, unless there is a contraindication to the mRNA vaccines, or as advised by the Medical Health Officer or an allergist,⁷ due to infrequent (1:50,000) but serious Vaccine-Induced Thrombotic Thrombocytopenia (VITT) blood clotting events after the first dose.¹⁴ The risk of VITT is more than six times lower for the second dose (1:600,000). People who had the VAXZEVRIA (AstraZeneca) vaccine for their first dose have the option of receiving VAXZEVRIA (AstraZeneca) for their second dose or receiving an mRNA vaccine as their second dose. Receiving a mixed vaccine series (VAXZEVRIA (AstraZeneca) for first dose and an mRNA vaccine for the second dose) is permitted based on small studies that suggest that this is likely safe and likely as effective and may be even more effective, but not enough is known to make firm conclusions and data collection is ongoing. There may also be heightened side effects experienced with a mixed vaccine series.

The BCCDC has prepared two information sheets to help navigate that choice:

- For health care professionals: www.bccdc.ca/Health-Info-Site/Documents/COVID-19_vaccine/Doctor_letter_Recommendations_AZ_COVISHIELD.pdf)
- **For patients:** Why your 2nd dose is important (http://www.bccdc.ca/Health-Info-Site/Documents/COVID-19_vaccine/2ndDose_Importance.pdf)

The Janssen COVID-19 Vaccine (Janssen)⁵ one-dose viral vector vaccine is now available in limited supply in B.C. However, mRNA vaccines are preferred over viral vector vaccines due to better effectiveness and immunogenicity of mRNA vaccines and the possible adverse effects specifically associated with viral vector vaccines (e.g., Thrombosis and Thrombocytopenia Syndrome [TTS]).¹⁴



The Novavax Nuvaxoid COVID-19 vaccine was authorized for use on February 17, 2022 by Health Canada. Novavax is a different class of vaccination, a protein subunit vaccine, that will give British Columbians another option to protect themselves against COVID-19. This vaccine is available to people aged 18 years and older. It is a two-dose vaccine and a limited number of doses will be available in B.C.¹⁵

There is still uncertainty as to whether the currently available COVID-19 vaccines are efficacious in adults and children with cancer or undergoing therapy for their cancer (cytotoxic chemotherapy, endocrine therapy, targeted therapy, immunotherapy) and/or radiation therapy (external-beam, brachytherapy, or systemic), as well as to the timing of immunization in relation to their cancer treatments.

Is the COVID-19 vaccine recommended for people with hematologic malignancies and HSCT and/or CAR-T recipients?

The risk of mortality from COVID-19 disease is higher in patients with cancer, including patients with hematologic malignancies and HSCT recipients.¹⁶⁻¹⁹ One study found that more severe forms of COVID-19 disease, including those requiring ICU admission, were more frequent in patients with hematologic malignancies hospitalized with COVID-19, and led to mortality nearly four times higher than that of the general population with COVID-19 and 41 times higher than that of hematologic malignancy patients without COVID-19.¹

COVID-19 vaccine vaccines are not contraindicated and should be encouraged for people with hematologic malignancies and HSCT and/or CAR-T recipients and as per BC Public Health recommendation for age eligibility. This recommendation is based on the following:

- The National Advisory Committee on Immunization (NACI) recommends that immunosuppressed individuals be offered the vaccine if the benefits of vaccines outweigh the potential risks.⁷
- Patients with blood cancer have an increased risk of death related to COVID-19 infection.^{1,16-19}
- The United Kingdom, the United States, France, and Australia have prioritized patients with cancer for COVID-19 vaccinations, highlighting the high COVID-19 risk faced by these patients.²⁰

Is the COVID-19 vaccine efficacious and safe in people with hematologic malignancy patients and HSCT and/or CAR-T recipients?

There is still uncertainty about the efficacy of COVID-19 vaccines in patients with blood cancer and/or have undergone HSCT or CAR-T cell therapy in the last six months. As with most vaccines, there is a potential for diminished immune response in individuals who are immunocompromised due to their disease or treatment. In addition, patients with active cancer or undergoing active cancer treatment seemed to be generally excluded from the COVID-19 vaccine trials. However, in the COMIRNATY (Pfizer-BioNTech) vaccine trial, 3.9% of enrolled participants had a malignancy.²¹

There are currently no known factors that would predispose these individuals to adverse events associated with the vaccines.²⁻⁵ At the time of authorization, there are no known serious warnings or precautions related to the vaccines in patients with cancer.²⁻⁵

Clinical Guidance on COVID-19 Vaccines for People with hematological malignancies and those who have undergone hematopoietic stem cell transplant or CAR-T

3



Small exploratory studies have shown lower antibody responses in patients with advanced cancer and haematological malignancies²¹ following vaccination compared to controls. However, it is unclear how much antibody is needed for protection and/or the role of other immunological responses.

There is one study that suggests that a third dose of COVID-19 vaccine in immunocompromised patients can increase antibody levels.²² Small studies on third doses of the mRNA COVID-19 vaccines have shown that immunogenicity (immunity measured in the blood) may increase with a third dose. The safety of a third dose is unknown at this time, but in these small studies reactions were found to be similar to that of prior doses. The impact of additional doses on the worsening of underlying disease or on rare adverse events, including the risk of myocarditis and/or pericarditis, is unknown at this time.¹¹

The frequency and severity of adverse events following vaccination with an mRNA COVID-19 vaccine in these populations were comparable to that of non-immunosuppressed individuals in these studies and what was reported in clinical trials. Safety data in these populations following vaccination with a viral vector vaccine is not available.⁸

Informed consent should include discussion about the possibility that individuals who are immunosuppressed may have a diminished immune response to any of the authorized COVID-19 vaccines.¹¹

Observed short-term adverse effects with the mRNA-based COVID-19 vaccines have been similar to those seen with seasonal influenza vaccination. Still, they may be more pronounced after the second COVID-19 vaccine dose (e.g. injection site pain/erythema, fever, headache, fatigue, and myalgia/arthralgia).^{2-5, 23} Safety results in the allogeneic HSCT patient population seem comparable.²³ Any long-term side effects of COVID-19 vaccines are not yet known, but Health Canada continues to monitor any adverse events following vaccination through their post-authorization surveillance process.

Immunocompromised patient populations are diverse and the relative degree of immunodeficiency will depend on the underlying condition, the progression of the disease, and the type and timing of treatment received. Therefore, the balance of potential benefit and risk associated with COVID-19 vaccination should be assessed on an individual basis (Table 1).

Are there any specific contraindications or exceptions for those within the hematologic malignancy, HSCT and/or CAR-T recipient patient populations?

Blood counts

Patients with blood cancer and HSCT or CAR-T recipients may experience low blood counts, either due to their disease or treatment, which could impact individual decision-making around receipt of COVID-19 vaccinations and timing of vaccinations relative to their treatments. COVID-19 vaccination should be deferred in patients unwell with neutropenia until well,^{2,3} but may be considered in well patients with disease-related chronic neutropenia where neutrophil recovery is not expected.²⁴



Allergy

The above noted COVID-19 vaccines are contraindicated in individuals with a history of severe allergic reaction to any component of the vaccines, including non-medicinal ingredients such as polyethylene glycol (PEG) or polysorbate-80, or a history of anaphylaxis after administration of a previous dose of COVID-19 vaccine using a similar platform (mRNA or viral vector).⁷ People with a history of anaphylaxis without known or obvious cause, and those with suspected hypersensitivity or non-anaphylactic allergy to COVID-19 vaccine components, are advised to consult with an allergist prior to immunization.²⁵ BCCDC has a list of the individual components and their purpose in the vaccines (www.bccdc.ca/health-info/diseases-conditions/covid-19/covid-19-vaccine/vaccines-for-covid-19). For a complete list of components in the vaccine, consult the vaccine monographs found at:

- tozinameran (COMIRNATY, Pfizer BioNTech): <https://covid-vaccine.canada.ca/info/pdf/pfizer-biontech-covid-19-vaccine-pm1-en.pdf>
- elasmomeran (SPIKEVAX, Moderna): <https://covid-vaccine.canada.ca/info/pdf/covid-19-vaccine-moderna-pm-en.pdf>
- ChAdOx1-S (VAXZEVRIA, AstraZeneca): <https://covid-vaccine.canada.ca/info/pdf/astrazeneca-covid-19-vaccine-pm-en.pdf>
- Ad26.COVS.2.S (Janssen COVID-19 Vaccine, Janssen): <https://covid-vaccine.canada.ca/info/pdf/janssen-covid-19-vaccine-pm-en.pdf>

For individuals with a history of anaphylactic reaction to a previous dose of an mRNA COVID-19 vaccine, re-vaccination (i.e., administration of a subsequent dose in the series when indicated) may be offered with the same vaccine or the same mRNA platform if a risk assessment deems that the benefits outweigh the potential risks for the individual and if informed consent is provided. Prior to revaccination, consultation with an allergist or another appropriate physician (e.g., Medical Health Officer) is advised. If re-vaccination is going ahead, vaccine administration should be done in a controlled setting with expertise and equipment to manage anaphylaxis, with an extended period of observation of at least 30 minutes after re-vaccination.

Health Canada continues to monitor any adverse events following immunization through their post-authorization surveillance [process](#).

Other vaccines

Currently, it is recommended that COVID-19 vaccines can be given concomitantly with, or any time before or after any other indicated vaccine.²⁶⁻²⁸ This is a change from the previous recommendation for a 14-day interval before or after receipt of a COVID-19 vaccine. The original advice against co-administration was based on a cautionary approach, as specific studies of co-administration with other vaccines have not been performed. However, substantial data have now been collected regarding the safety of COVID-19 vaccines currently authorized by Health Canada. Extensive experience with non-COVID-19 vaccines has demonstrated that immunogenicity and adverse event profiles are generally similar when vaccines are administered simultaneously as when they are administered alone.

Clinical Guidance on COVID-19 Vaccines for People with hematological malignancies and those who have undergone hematopoietic stem cell transplant or CAR-T

5



Are there specific recommendations or considerations for safe and/or most effective administration?

1. Blood counts

Patients with blood cancer and HSCT or CAR-T recipients may have lowered blood counts related to the underlying disease or therapy. If blood counts (platelet count and neutrophil count) are low due to therapy and timing of recovery can be anticipated, e.g. 1 week prior to the next cyclical chemotherapy or maintenance cycle, the timing of vaccination should be scheduled accordingly (please see Table 1). However, where the timing of blood count recovery is unclear or not anticipated, e.g. marrow failure syndromes, then vaccination should not be delayed solely for this reason.

There is no consensus on an adequate platelet count for IM injections. Still, practical suggestions include using a platelet threshold of $>20 \times 10^9/L$, administering the vaccine after platelet transfusion if receiving regular transfusions, and applying firm pressure at the injection site for at least 5 minutes.²⁹

2. Anti-coagulant therapy

As per Thrombosis Canada recommendations,³⁰ anti-coagulation should not be a barrier for administering COVID-19 vaccination to patients on warfarin (INR monitoring not required prior to vaccination), novel oral anticoagulants (apixaban, dabigatran, edoxaban, rivaroxaban) or antiplatelet agents (aspirin, clopidogrel, ticagrelor). Patients on therapeutic dose low-molecular weight heparin (dalteparin, tinzaparin, enoxaparin, nadroparin) or fondaparinux may consider delaying their anti-coagulant dose on the day of vaccination until after the IM injection. For patients on any of the above, applying pressure to the injection site for 3 to 5 minutes post vaccination is recommended to reduce bruising.

3. Special considerations for immunotherapy

a. Therapies targeting B-cells including anti-CD20, CD19, CD22 targeting antibodies, or BiTEs: Patients receiving these agents may have a reduced immune response to vaccines in general that can extend to up to 6 months following treatment completion. If possible, patients should receive both doses of vaccine prior to starting these therapies. If patients are on, or have recently been treated with these agents, when they received the first 2 doses of vaccine, a 3rd dose is recommended to be administered at least 28 days after the 2nd dose, with consideration given to delaying to 3 months after therapy with B-cell directed therapies due to likelihood of impaired immune response.

b. Checkpoint inhibitors:

Previous studies have not signalled an increased risk of complications of COVID-19 for patients on checkpoint inhibitors such as CTLA-4 inhibitors (e.g., ipilimumab), PD-1 inhibitors (e.g., nivolumab, pembrolizumab) and PD-L1 inhibitors (e.g., atezolizumab, durvalumab). There have been theoretical concerns of an enhanced immune reaction, particularly with CTLA-4 inhibitors. However, given the seriousness of COVID-19 infection, vaccination is still recommended in this group even if a four-week window cannot be confirmed.

4. Timing of COVID-19 vaccines in relation to therapy

There are no known studies regarding the timing of COVID-19 vaccination in relation to therapy for blood cancer. The

Clinical Guidance on COVID-19 Vaccines for People with hematological malignancies and those who have undergone hematopoietic stem cell transplant or CAR-T

6



COMIRNATY (Pfizer-BioNTech), SPIKEVAX (Moderna) and VAXZEVRIA (AstraZeneca) vaccines are given as two injections with optimal protection assumed after the second dose for the general population.²⁻⁵ Efficacy and duration of immunity are continuously being evaluated and recommendations are evolving rapidly. Therefore, patients should follow current BCCDC guidance for the recommended number of and interval between COVID-19 vaccine doses.

In general, it is preferred that patients complete their COVID-19 vaccination series ideally 14 days prior to starting immunosuppressive therapy.³¹

***However, life-saving or prolonging therapy should not be delayed solely to complete vaccination.**

Recommendations for timing of COVID-19 vaccination for patients with hematologic malignancies (either completed, starting or already receiving treatment) and patients who have undergone HSCT or CAR-T cell therapy in the past 6 months are described in Table 1 below.

Any other timing should involve a case-by-case assessment based on:

- Risk of morbidity related to COVID-19 infection (including local incidence of the pandemic, cancer type, comorbidities that confer higher risk categories in general population, etc.),
- Cancer-related morbidity due to delay of active treatment, and
- Suboptimal immunity due to insufficient time window between vaccination and immunosuppressive therapy.

Table 1. Suggested timing of COVID-19 vaccination in patients with hematologic malignancies³²⁻³⁵

Therapy	Suggested timing of COVID-19 vaccine
Cyclical chemotherapy – <i>prior to starting</i> (including hypomethylating agents)	<ol style="list-style-type: none"> Ideally complete vaccination at least 2 weeks prior to starting* Alternatively, complete vaccination between cycles of therapy if clinically not appropriate to wait to complete vaccination
Cyclical chemotherapy – <i>between cycles</i> (including hypomethylating agents)	Give vaccine dose(s) between cycles: <ul style="list-style-type: none"> Upon count recovery (<i>if anticipated to recover</i>)** about 1 week prior to starting subsequent cycle <i>Note: Avoid on same day as treatment</i>
Single agent small molecule inhibitors (e.g. kinase inhibitors or continuous oral chemotherapy, BTK inhibitors)	No specific timing
Immunomodulatory agents	Avoid on same day as treatment
Proteasome inhibitors (e.g. bortezomib)	Avoid on same day as treatment

Clinical Guidance on COVID-19 Vaccines for People with hematological malignancies and those who have undergone hematopoietic stem cell transplant or CAR-T



Therapy	Suggested timing of COVID-19 vaccine
Check point inhibitors	Avoid on same day as treatment
CD19, CD20, CD22 targeted therapy (e.g. monoclonal antibodies)	No specific timing [†]
Other monoclonal antibodies	No specific timing
Systemic Corticosteroids	Cyclical corticosteroids as part of chemotherapy regimens – ideally vaccinate on days when not receiving corticosteroids Continuous corticosteroids – no specific timing ***
Autologous HSCT [§] [¥]	Pre-HSCT: ≥ 2 weeks prior to starting conditioning chemotherapy Post-HSCT: > 3 months post-HSCT
Allogeneic HSCT [§] [¥]	Pre-HSCT: ≥ 2 weeks prior to starting conditioning chemotherapy Post-HSCT: > 3 months post-HSCT [†]
CAR-T cell therapy [§]	Pre-CAR-T cell therapy: ≥ 2 weeks prior to starting lymphodepleting chemotherapy Post-CAR-T cell therapy: > 3 months post-CAR-T cell therapy
Aplastic Anemia	Not on therapy or completed therapy with counts in acceptable range: No specific timing required Post-therapy: > 3 months post-initiation of cyclosporine/ATG [†]
Intravenous immunoglobulin (IVIg) - Not COVID-19 specific	No specific timing
Under Observation - Not scheduled for therapy OR completed planned therapy	No specific timing

* In general, it is preferable to complete vaccination before starting immunosuppressive therapy if possible (based on timing of therapy and vaccine availability). However, life-saving or prolonging therapy should not be delayed solely to complete vaccination. Some immunity may be achieved following the first dose of the vaccine.

** Some patients may not have adequate counts either prior to or between cycles of therapy. The benefit likely outweighs the risk, and these patients should proceed to vaccination regardless of neutrophil count and with platelet transfusion support if required.

*** Ideally high dose systemic corticosteroids (> 0.5 mg/kg/day prednisone or equivalent) should be avoided or completed 28 days prior to vaccination; if this is not possible, proceed with vaccination.

[†] Due to likelihood of impaired immune response to vaccination within 3 months of receiving B-cell directed monoclonal antibodies and ATG, consider delaying to 3 months post-therapy.

Clinical Guidance on COVID-19 Vaccines for People with hematological malignancies and those who have undergone hematopoietic stem cell transplant or CAR-T



[§] Rationale for consideration of delaying COVID-19 vaccination for > 3 months after HSCT and CAR-T cell therapy includes:

- Vaccine response is expected to be sub-optimal;
- Antibody testing cannot be evaluated as standard of practice;
- Revaccination post-HSCT and CAR-T cell therapy is recommended. The attestation form for “Revaccination following Hematopoietic Stem Cell Transplant” can be obtained from transplant physicians.

[¥] If local COVID-19 transmission rates are high, consider prioritization of COVID-19 vaccination. Routine post-HSCT vaccinations may be given at the same time as the COVID-19 vaccines but may be delayed at the discretion of the patient or medical professional.³²

References

1. Passamonti *et al.* Clinical characteristics and risk factors associated with COVID-19 severity in patients with haematological malignancies in Italy: a retrospective, multicentre, cohort study. *Lancet Haematol.* 2020;7(10):e737-45.
2. Pfizer-BioNTech. COMIRNATY product monograph. Kirkland, Quebec. 19 November 2021.
3. Moderna. SPIKEVAX product monograph. Oakville, Ontario. 12 November 2021.
4. AstraZeneca. VAXZEVRIA product monograph. Mississauga, Ontario. 19 November 2021.
5. Janssen. Janssen COVID-19 vaccine product monograph. Toronto, Ontario. 23 November 2021.
6. National Advisory Committee on Immunization. Recommendation on the use of the Pfizer-BioNTech COVID-19 vaccine (10 mcg) in children 5-11 years of age. 19 November 2021. Available at: <https://www.canada.ca/content/dam/phac-aspc/documents/services/immunization/national-advisory-committee-on-immunization-naci/recommendations-use-covid-19-vaccines/pfizer-biontech-10-mcg-children-5-11-years-age/pfizer-biontech-10-mcg-children-5-11-years-age.pdf>
7. National Advisory Committee on Immunization. Recommendations on the use of COVID-19 vaccine(s). 22 July 2021. Available at: <https://www.canada.ca/en/public-health/services/immunization/national-advisory-committee-on-immunization-naci/recommendations-use-covid-19-vaccines.html> Accessed on: 20 September 2021.
8. BC Centre for Disease Control. COVID-19 Vaccine. Getting a Vaccine. Getting your first and second doses. Updated: 28 February 2022. Available at <http://www.bccdc.ca/health-info/diseases-conditions/covid-19/covid-19-vaccine/getting-a-vaccine#first> Accessed on: 1 March 2022
9. Province of British Columbia. Get your booster dose. Youth 12 to 17. Clinically extremely vulnerable criteria for youth. <https://www2.gov.bc.ca/gov/content/covid-19/vaccine/booster#youth> Accessed: 24 February 2022.
10. BC Centre for Disease Control. COVID-19 Vaccine. Getting a Vaccine. Booster dose. Updated March 1 2022. Available at: <http://www.bccdc.ca/health-info/diseases-conditions/covid-19/covid-19-vaccine/getting-a-vaccine#booster> Accessed on: 1 March 2022.
11. National Advisory Committee on Immunization (NACI) rapid response: Additional dose of COVID-19 vaccine in immunocompromised individuals following 1- or 2- dose primary series. Sept 10 2021. <https://www.canada.ca/en/public-health/services/immunization/national-advisory-committee-on-immunization-naci/statement-september-10-2021-additional-dose-covid-19-vaccine-immunocompromised-following-1-2-dose-series.html>. Accessed on: Sept. 10 2021



12. BC Immunization Manual Update – Administrative Circular. Feb 3, 2022. http://www.bccdc.ca/resource-gallery/Documents/Guidelines%20and%20Forms/Guidelines%20and%20Manuals/Epid/CD%20Manual/Admin%20Circulars/2022/AC_2022-06_Chapter_2_Immunization_Feb_3_2022.pdf Accessed 5 February 2022.
13. BC Immunization Manual Update – Administrative Circular. September 14 2021. http://www.bccdc.ca/resource-gallery/Documents/Guidelines%20and%20Forms/Guidelines%20and%20Manuals/Epid/CD%20Manual/Admin%20Circulars/2021/AC_2021-35_Chapter_2_Immunization_Sep_14%20_2021.pdf. Accessed September 15, 2021.
14. Andrea Greinacher et al. The New England Journal of Medicine. Thrombotic Thrombocytopenia after ChAdOx1 nCov-19 Vaccination. DOI: 10.1056/NEJMoa2104840
15. Government of Canada. Novavax Nuvaxovid COVID-19 vaccine. <https://www.canada.ca/en/health-canada/services/drugs-health-products/covid19-industry/drugs-vaccines-treatments/vaccines/novavax.html> Accessed on: 1 March 2022.
16. Clift A, Keogh R, Diaz-Ordaz K et al. Living risk prediction algorithm (QCOVID) for risk of hospital admission and mortality from coronavirus 19 in adults: national derivation and validation cohort study. *BMJ*. 2020;371:m3731.
17. Wood *et al*. Outcomes of patients with hematologic malignancies and COVID-19: a report from the ASH Research Collaborative Data Hub. *Blood Adv*. 2020;4(23):5966-75.
18. Sharma *et al*. Clinical characteristics and outcomes of COVID-19 in haematopoietic stem-cell transplantation recipients: an observational cohort study. *Lancet Haematol*. 2021 Jan 19(8):e185-93.
19. Vijenthira *et al*. Outcomes of patients with hematologic malignancies and COVID-19: a systematic review and meta-analysis of 3377 patients. *Blood*. 2020;136(25):2881-92.
20. CADTH. COVID-19 mRNA vaccines for people with cancer. 22 December 2020 [accessed 23 December 2020]. https://cadth.ca/sites/default/files/covid-19/COVID-19_mRNA_vaccines_for_people_with_cancer.pdf
21. Monin L, et al. Safety and immunogenicity of one versus two doses of the COVID-19 vaccine BNT162b2 for patients with cancer: interim analysis of a prospective observational study. *Lancet Oncol*. 2021 Jun;22(6):765-778. doi: 10.1016/S1470-2045(21)00213-8. Epub 2021 Apr 27. PMID: 33930323; PMCID: PMC8078907.
22. Marlet, J., Gatault, P., Maakaroun, Z., Longuet, H., Stefic, K., Handala, L., Eymieux, S., Gyan, E., Dartigeas, C., & Gaudy-Graffin, C. (2021). Antibody Responses after a Third Dose of COVID-19 Vaccine in Kidney Transplant Recipients and Patients Treated for Chronic Lymphocytic Leukemia. *Vaccines*, 9(10), 1055. <https://doi.org/10.3390/vaccines9101055>
23. Polack FP *et al*. Safety and efficacy of the BNT162b2 mRNA Covid-19 vaccine. Supplementary appendix. *N Engl J Med* [published online ahead of print, 2020 Dec 10]. https://www.nejm.org/doi/suppl/10.1056/NEJMoa2034577/suppl_file/nejmoa2034577_appendix.pdf
24. Ali H, Ngo D, Aribi A, Arslan S, Dadwal S, Marcucci G, Nakamura R, Forman SJ, Chen J, Al Malki MM. Safety and Tolerability of SARS-CoV2 Emergency-Use Authorized Vaccines for Allogeneic Hematopoietic Stem Cell Transplant Recipients. *Transplant Cell Ther*. 2021 Jul 15:S2666-6367(21)01073-3. doi: 10.1016/j.jtct.2021.07.008. Epub ahead of print. PMID: 34274492; PMCID: PMC8280601.
25. BC Centre for Disease Control. COVID-19 Vaccine. Vaccine Considerations. Updated: 11 February 2022. Available at: <http://www.bccdc.ca/health-info/diseases-conditions/covid-19/covid-19-vaccine/vaccine-considerations> Accessed on: 1 March 2022



26. BC Centre for Disease Control. Communicable Disease Control Manual. Chapter 2: Immunization. Part 4 – Biological Products. 25 February 2022. http://www.bccdc.ca/resource-gallery/Documents/Guidelines%20and%20Forms/Guidelines%20and%20Manuals/Epid/CD%20Manual/Chapter%20%20-%20Imms/Part4/COVID-19_mRNA_Vaccine_BNT162b2_Pfizer-BioNTech.pdf Accessed on: 1 March 2022.
27. BC Centre for Disease Control. Communicable Disease Control Manual. Chapter 2: Immunization Part 4 - Biological Products. COVID-19 mRNA Vaccine BNT162b2. June 3, 2021 [accessed 3 June 2021]. http://www.bccdc.ca/resource-gallery/Documents/Guidelines%20and%20Forms/Guidelines%20and%20Manuals/Epid/CD%20Manual/Chapter%20%20-%20Imms/Part4/COVID-19_mRNA_Vaccine_BNT162b2_Pfizer-BioNTech.pdf
28. BC Centre for Disease Control. Communicable Disease Control Manual. Chapter 2: Immunization Part 4 - Biological Products. COVID-19 mRNA Vaccine mRNA-1273. June 3, 2021 [accessed 3 June, 2021]. http://www.bccdc.ca/resource-gallery/Documents/Guidelines%20and%20Forms/Guidelines%20and%20Manuals/Epid/CD%20Manual/Chapter%20%20-%20Imms/Part4/COVID-19_mRNA_Vaccine_mRNA-1273_Moderna.pdf
29. ASH-ASTCT COVID-19 and Vaccines: Frequently Asked Questions (Version 2.1; last updated December 23, 2020 [accessed 15 January 2021]). Input from Drs. Jeff Auletta, Roy Chemaly, Fareed Khawaja, Genovefa Papanicolaou, Josh Hill, Julie Kanter, Alpana Waghmare, Adrian Wiestner, John Wingard. <https://www.hematology.org/covid-19/ash-astct-covid-19-and-vaccines>
30. Thrombosis Canada Guidance on COVID-19 Vaccines and Anticoagulation. December 22, 2020 [accessed 19 January 2021]. <https://thrombosiscanada.ca/thrombosis-canada-guidance-on-covid-19-vaccines-and-anticoagulation/>
31. UK Chemotherapy Board. Clinician Frequently Asked Questions (FAQs) and guidance on COVID-19 vaccine for patients receiving Systemic Anti-Cancer Therapy. Version 1.0 Dec 2020 [accessed 12 January 2021]. https://b-s-h.org.uk/media/19241/clinician-faqs-and-guidance-on-covid19-vaccine-for-patients-receiving-sa_.pdf
32. Public Health England Guidance. COVID-19: the green book, chapter 14a. 11 December 2020 [accessed 15 December 2020]. <https://www.gov.uk/government/publications/covid-19-the-green-book-chapter-14a>
33. BSBMT&CT recommendations for the management of adult patients and allogeneic donors during the COVID-19 (causative agent the SARS-CoV-2 virus) outbreak. Version 5; 4 December 2020 [accessed 15 January 2021]. https://bsbmtct.org/wp-content/uploads/2020/12/BSBMTCT-COVID-19-Guidelines-5.0-Dec-2020_final.pdf
34. Centers for Disease Control and Prevention. Interim Clinical Considerations for Use of mRNA COVID-19 Vaccines Currently Authorized in the United States. [accessed 27 January 2021]. <https://www.cdc.gov/vaccines/covid-19/info-by-product/clinical-considerations.html>
35. British Society for Haematology. COVID-19 Vaccine in patients with haematological disorders. V1 21 December 2020 [accessed 19 January 2021]. <https://b-s-h.org.uk/media/19195/haematology-covid-19-v10-vaccination-statement-231220.pdf>

Authors

Dr Kevin Song, MD, FRCPC; Medical Director, Leukemia/Bone Marrow Transplant Program of British Columbia, Vancouver General Hospital

Clinical Guidance on COVID-19 Vaccines for People with hematological malignancies and those who have undergone hematopoietic stem cell transplant or CAR-T

11



Dr Sujaatha Narayanan, MBBS, MRCP, FRCPath; Leukemia/Bone Marrow Transplant Program of British Columbia, Vancouver General Hospital

Carmen Mountford, BSc, BScPharm, ACPR; Clinical Pharmacist, Leukemia/Bone Marrow Transplant Program of British Columbia, Vancouver General Hospital, Lower Mainland Pharmacy Services

Reviewers

Dr Caron Strahlendorf MB, FRCPC, FCP; Clinical Associate Professor, Division of Hematology and Oncology, Department of Pediatrics, Faculty of Medicine, University of British Columbia

Dr Alissa Wright, BSc, MD, FRCPC, MSc; Head, Division of Infectious Diseases, Vancouver General Hospital

Dr Helen Anderson, Program Medical Director, Systemic Therapy, BC Cancer

Dr Stephen Nantel, Head, Division of Hematology, Leukemia/Bone Marrow Transplant Program of British Columbia, Vancouver General Hospital

Dr Paul Yenson, Medical Lead, General Hematology, Vancouver General Hospital

Dr Maryse Power, Associate Medical Director, Leukemia/Bone Marrow Transplant Program of British Columbia, Vancouver General Hospital

Dr Kerry Savage, Medical Oncologist, BC Cancer

Dr Laurie Sehn, Lymphoma Tumor Group Chair, BC Cancer

Nursing leadership, Leukemia/BMT Program of BC, Vancouver General Hospital

