

Coronavirus COVID-19

BC Centre for Disease Control | BC Ministry of Health



COVID-19 Infection Prevention and Control: Guidance for Community-Based Physicians, Nursing Professionals and Midwives in Clinic Settings

June 4, 2021

Contents

- A. Introduction 2**
 - 1. COVID-19 Immunization 2
- B. Infection Prevention and Control Measures 2**
 - Public Health Measures 3**
 - 1. Public Health Order – Workplace Safety Plans 5
 - 2. Public Health Order - Gatherings and Events 5
 - 3. Case Finding, Contact Tracing and Outbreak Management 5
 - 4. Self-isolation and Quarantine 4
 - Environmental Measures 4**
 - 5. Cleaning and Disinfection 4
 - 6. Physical Changes to the Clinic 5
 - Administrative Measures 7**
 - 7. Staff Screening 7
 - 8. Patient Screening and Management 6
 - 9. Physical Distancing and Minimizing Physical Contact 8
 - Personal Measures 9**
 - 10. Stay Home When Sick 9
 - 11. Routine Practices 9
 - Guidance for Personal Protective Equipment (PPE) 10**
 - 12. Key Personal Protective Equipment Recommendations 10
 - 13. PPE Guidance for Patients and Visitors 12
 - Office Management 12**
 - 14. Clinic Response Planning and Organization 12
 - 15. Staff Education and Communication 12
 - 16. Psychosocial Support 12
 - 17. Staff Scheduling and Reassignments 12
 - 18. Sick Leave Policy 13
 - Key Resources 12**
 - References 15**



Ministry of Health



BC Centre for Disease Control

If you have fever, a new cough, or are having difficulty breathing, call 8-1-1.



Appendix A: COVID-19 Community Clinic Care Preparedness Checklist15

A. Introduction

This document provides a practical guide for primary care practitioners, community-based physicians, nursing professionals, midwives and related staff in clinics and urgent and primary care centres to support appropriate office-based infection prevention and control (IPC) practices to mitigate the impact of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), the causative agent of COVID-19.

This guidance outlines the IPC measures to provide care safely in these settings, including patients with confirmed COVID-19 and suspected COVID-19 cases (e.g., having symptoms of COVID-19 or risk factors (required to self-isolate following close contact or travel)). It is based on the latest available best practice and scientific evidence. The guidance may change as new information becomes available. For COVID-19 variants of concern, recommendations for IPC measures remain the same and should be strictly followed and reinforced. See [Guidance on SARS-CoV-2 Variants of Concern](#) for more information.

Please see [appendix A](#) for a COVID-19 community clinic care preparedness checklist to assist with implementing this guidance.

COVID-19 Immunization

Overall, approved COVID-19 vaccines in Canada are effective against COVID-19. We are continuing to learn about the impact that immunization has on SARS-CoV-2 transmission, and their effectiveness against certain variants of concern. As the evidence evolves, public health and IPC guidance for individuals who have received their COVID-19 immunizations will be updated as needed.

Some of the side effects from COVID-19 vaccines are similar to the common symptoms of COVID-19. Individuals experiencing any symptoms of COVID-19 after receiving their immunizations are to continue to use the [BC COVID-19 Self-Assessment tool](#) to determine if testing for COVID-19 is required.

Currently in health-care facilities, regardless of whether an individual (e.g., patient, health-care worker, visitor) has received a COVID-19 vaccine(s) and when they received it, they must continue to follow local processes for COVID-19 screening and managing COVID-19 like symptoms. When providing care to symptomatic patients, health-care workers (HCWs) must continue to conduct point-of-care risk assessments (PCRAs) and implement additional precautions as needed to prevent the transmission of SARS-CoV-2.

For further information, please see the following resources:

[NACI Recommendations on the use of COVID-19 vaccines](#)

[BCCDC Monitoring vaccine update, safety and effectiveness \(March 30, 2021\)](#)


[BCCDC Getting a vaccine](#)

B. Infection Prevention and Control Measures

Implementation of IPC measures helps create a safe environment for health-care workers, staff and patients. A hierarchy of IPC measures for communicable disease describes the measures that can be taken to reduce the transmission of COVID-19. IPC measures at the top are more effective and

protective than those at the bottom. By implementing a combination of measures at each level, the risk of COVID-19 is substantially reduced.

Hierarchy for Infection Prevention and Exposure Control Measures for Communicable Disease



Public health measures are society-wide actions to limit the spread of the SARS-CoV-2 virus and reduce the impact of COVID-19. The provincial health officer has implemented public health orders, including prohibiting mass gatherings, requiring travellers to self-isolate (or quarantine upon arrival in B.C. according to the federal quarantine order), effective testing, case finding and contact tracing, and emphasizing the need for people to stay home when they are sick.

Environmental measures are physical changes in a setting that reduce the risk of exposure by isolation or ventilation. Examples include being in outdoor spaces, having suitable ventilation and air exchange, using visual cues for maintaining physical distance, erecting physical barriers where appropriate, and frequent cleaning and disinfection of work and living spaces.

Administrative measures are the implementation of policies, procedures, training, and education. Examples of these include decreased density of staff and patients in clinics, staggered appointments, and using virtual health where appropriate.

Personal measures are actions individuals can take to both protect themselves, as well as to prevent the spread to others. Examples include washing hands frequently, coughing into an elbow, and staying home from work when sick.

Personal protective equipment (PPE) is the last and least effective of the infection prevention and exposure control measures. It is not effective as a stand-alone preventive measure and should only be

considered after implementing all other measures. PPE must be suited to the task and must be worn and disposed of properly.

Clinics can implement a combination of measures from the different levels described above to protect against COVID-19 as detailed below.

Public Health Measures

1. [Public Health Order – Workplace Safety Plans](#)

By order of the Provincial Health Officer (PHO), all employers in B.C. are required to post a copy of a [WorkSafe BC COVID-19 safety plan](#) that outlines the policies, guidelines and procedures employers have put in place to reduce the risk of COVID-19 transmission.

2. [Public Health Order - Gatherings and Events](#)

By order of the PHO, restrictions are in place for in-person events and community-based gatherings. Please see the [province-wide restrictions](#) for more information

3. Case Finding, [Contact Tracing](#) and [Outbreak Management](#)

When a COVID-19 positive person is identified by public health staff, significant efforts are undertaken to determine if that person is part of a cluster of cases or a local outbreak. Specific public health measures are implemented in facilities where an outbreak occurs to prevent further transmission of COVID-19 and keep others safe in the workplace.

Operators should maintain an updated list of **staff and patients' first and last names, their phone numbers or emails, and the dates of clinic attendance**, to support public health staff in potential contact tracing efforts.

4. [Self-isolation](#) and Quarantine

HCWs and staff with common cold, influenza, or symptoms of COVID-19 should stay home, get assessed by their health-care provider and get tested for COVID-19 if their symptoms warrant a test in accordance with [provincial COVID-19 testing criteria](#).

When someone has symptoms of COVID-19, they should self-isolate and follow directions provided by their health-care provider.

Self-isolation is also advised for those who are considered a close-contact of a confirmed case and are waiting to see if they develop COVID-19 illness in accordance with public health direction.

Quarantine is a term typically reserved for people who return from travel outside the country and are at risk of developing COVID-19 as outlined under the federal [quarantine order](#).

Environmental Measures

5. [Cleaning and Disinfection](#)

Regular cleaning and disinfection are essential for preventing the transmission of COVID-19 from contaminated objects and surfaces.

Clinic spaces should be cleaned and disinfected in accordance with BC Centre for Disease Control's (BCCDC) [environmental cleaning and disinfectants for clinic settings](#) document.

This includes the following:

- Minimize equipment brought into the exam room.
- Clean and disinfect shared reusable equipment **between patients**.
 - This includes stethoscopes, blood pressure cuffs, otoscopes, baby scales, tables and examination beds.
- Clean and disinfect frequently touched surfaces at least **twice a day**.
 - These include medical equipment, doorknobs, light switches, telephones, keyboards, mice, pens, clipboards or plastic binding on charts, cell phones, toys and all hard surfaces in bathrooms (e.g., sinks, faucets, handles).
- Establish clear roles and responsibilities for those responsible for cleaning and disinfecting each piece of medical or non-medical equipment, and environmental surfaces.
- Clean and disinfect the procedure and examination rooms at least **once a day**.
- Clean and disinfect all surfaces that are visibly dirty.
- Use a hospital grade disinfectant with a drug identification number, in accordance with manufacturer's instructions, and contact (wet) time requirements to ensure pathogens have been killed.
- Limit items that are not easily cleaned (e.g., fabric or soft items).
- Follow routine procedures for waste disposal and laundry management.
- Empty garbage containers daily.
- Wear disposable gloves when cleaning blood or body fluids (e.g., runny nose, vomit, stool, urine).
- Perform hand hygiene before wearing and after removing gloves.

The risk of transmission of COVID-19 via paper or other paper-based products is low^{1,2}. As such, there is no need to limit the distribution of paper resources, such as leaflets, to patients because of COVID-19.

6. Physical Changes to the Clinic

Facilities may not be able to adopt all of the following measures; however, consideration should be given to incorporating **as many measures as possible**:

- Spread people out into different areas or have patients wait outside the clinic until their appointment time. Consider different configurations to allow distance between staff and patients, and between patients. (e.g., different chair positioning in waiting rooms).
- Create designated COVID-19 triage, waiting areas and examination rooms, if possible.
- Provide alcohol-based hand rubs (ABHR) with a minimum of 70% alcohol at the reception counter and near exam room doors, if available. A higher concentration of alcohol is required to eliminate a wide range of microorganisms, including COVID-19³.
- Exam rooms:
 - The exam room(s) closest to the entrance should be designated for patients with respiratory symptoms in order to allow rapid isolation pending formal assessment.
 - Exam rooms should be emptied of all but the bare minimum equipment (e.g., exam table, one chair, blood pressure cuff, lights).
 - Minimize sterile and clean supplies located in exam rooms. Keep supplies in closed cabinets/containers to minimize the risk of contamination.
 - Set up a PPE station outside of exam rooms.

- Keep frequently used doors open (where appropriate) to avoid recurrent contamination of doorknobs and high-touch contact points.
- Place signage in the clinic. Up-to-date COVID-19 signage and information is available for download from the [BCCDC signage and posters webpage](#) (available in multiple languages). Post appropriate posters and signage at entrance/exit doors, the reception area and all exam rooms. Post hand hygiene signage near all sinks.
- Heating, ventilation and air conditioning (HVAC) systems should be properly installed and regularly inspected and maintained according to HVAC standards by the Canadian Standards Association and other building code requirements. Where feasible, optimize HVAC systems in patient care areas and rooms, especially where patients suspected or confirmed of having COVID-19 are accommodated and cared for. Information on optimizing HVAC systems in the context of COVID-19 can be sought from [WorkSafeBC](#), the [American Society of Heating, Refrigerating and Air-Conditioning Engineers](#), [Alberta Health Services COVID-19 Scientific Advisory Group](#), [Public Health Ontario](#). When adjustments are needed, it would be beneficial to solicit the expertise of HVAC specialists.
- Where physical distancing cannot be maintained, consider physical barriers⁴ such as:
 - Installing clear partitions such as plexiglass or plastic film, at reception counters and other areas. Change or clean and disinfect clear partitions frequently⁴.
 - Installing dividers (e.g., free-standing partitions or privacy curtains) between patient stations. Change or clean and disinfect dividers frequently⁴.
 - For further information about laundry management see the BCCDC website for an [information sheet for environmental service providers in health-care settings](#).
 - If privacy curtains are used in the facility, remove and launder them when visibly soiled, at discharge if the patient is on additional precautions, and at least quarterly⁵.

Administrative Measures

7. Staff Screening

Before each shift, each health-care worker (HCW) must self-screen for COVID-19 symptoms in accordance with current orders from the PHO and the employer's COVID-19 safety plans and follow measures outlined in the [COVID-19 health-care worker self-check and safety checklist](#).

8. Patient Screening and Management

Before each appointment, every patient should be screened for COVID-19 risk using the [COVID-19 Patient Screening Tool for Direct Care Interactions](#).

Consideration should be given to the management of patients with suspected or confirmed COVID-19 cases in the clinic. Options include having dedicated areas for COVID-19 patients and/or scheduling appointments for these patients later in the day; as well as seeing high-risk patients (e.g., elderly and those with chronic illnesses) as the first appointments of the day.

For clinics seeing patients on a walk-in basis, screen patients for COVID-19 risk upon entry to the clinic using the [COVID-19 entrance screening tool for health-care facilities](#).

Plan for patient scheduling, patient flow and triage. Avoid non-essential visitors accompanying patients, where possible.

Explore collaborative approaches to clinic management and patient care. This could consist of two or more health-care practices located in the same building developing ways to work together, with one practice having virtual-only appointments and the other having in-person appointments. Any change in practice should be clearly communicated to patients and staff.

Pre-Visit Messaging

Office telephone message/voice mail and the health-care practice website(s) should clearly instruct patients where to seek up-to-date instructions on assessment for COVID-19. For assessment, patients can be directed to the [BC COVID-19 Self-Assessment Tool](#) and BCCDC's website for information on [lab testing](#).

Pre-Appointment Triage

A comprehensive triage process starts prior to a patient arriving at the clinic. When booking a patient's appointment:

Ask the COVID-19 screening questions (see BCCDC's website for a sample [COVID-19 patient screening tool for direct care interactions & routine point-of-care risk assessment tool](#)), including whether the patient has symptoms of COVID-19 or has been confirmed to have COVID-19.

Ensure the patient is informed of the recommended precautions and requirements such as wearing a medical mask, hand hygiene, respiratory hygiene and environmental cleaning and disinfection.

9. Physical Distancing and Minimizing Physical Contact

Physical distancing means maintaining a distance of two metres between people. The following physical distancing strategies should be implemented where possible:

- Reduce the number of people in waiting rooms.
- Avoid close greetings (e.g., handshakes).
- Stagger appointment times.
- Stagger break times for staff.
- Manage the flow of people in common areas, including waiting rooms and hallways.
- Minimize the number of caregivers and other individuals who are not patients entering the clinic, as much as is practical to do so.
- Post signs and remind everyone to practice diligent hand hygiene and maintain physical distance while they are in the clinic.

Personal Measures

10. Stay Home When Sick

All staff must stay home and self-isolate:

- who have symptoms of COVID-19 OR
- have travelled outside Canada in the last 14 days OR
- have been identified as a close contact of a confirmed COVID-19 case.

To determine the need for testing and whether to self-isolate, direct staff to the [BC COVID-19 self-assessment tool](#).

If concerned, staff can be advised to contact 8-1-1, their health-care provider or the local public health unit to seek further advice.

See the BCCDC's [exposures and return to work for health-care workers](#) guidance on HCWs exposed to COVID-19 while at work, what to do if a staff member becomes ill, how long to self-isolate and criteria for return to work for those with symptoms.

11. Routine Practices

Routine practices and additional precautions should be in place at all times in all health-care settings. This includes performing diligent hand hygiene, adhering to respiratory hygiene, conducting a PCRA, and appropriate use of PPE.

Hand Hygiene

Rigorous hand hygiene with plain soap and water or ABHR is the most effective way to reduce the spread of illness.

Wash hands with plain soap and water for at least 20 seconds. Antibacterial soap is not needed for COVID-19. If sinks are not available, use ABHR containing at least 70% alcohol.

If hands are visibly soiled, ABHR may not be effective at eliminating respiratory viruses. Soap and water are preferred when hands are visibly dirty.

To learn about how to perform hand hygiene, please refer to BCCDC's website for a [hand hygiene poster](#).

Strategies to support and promote diligent hand hygiene:

- Hand hygiene stations should be set up at the clinic entrance, so everyone can perform hand hygiene when they enter.
- Hand hygiene sinks, plain soap dispensers, paper towel holders, hand sanitizer dispensers, and related supplies, should be readily available throughout the facility.
- Post signs or posters to promote regular hand hygiene.
- Paper towels should be disposed of in non-touch wastebaskets lined with a garbage bag.

For patients, HCWs and staff, hand hygiene must be performed:

- On entering the clinic;
- On entering the examination room;
- On leaving the examination room;
- After using the washroom;

- After using a tissue for their face; and
- After coughing or sneezing.

For HCWs and staff, hand hygiene must also be performed:

- Before and after contact with patient or the patient care environment;
- Before and after breaks;
- Before clean or sterile procedures;
- After risk of body fluid exposure;
- Before putting on/donning PPE; and
- In between each step when taking off/doffing PPE.

Respiratory Etiquette

Respiratory etiquette is also known as respiratory and cough hygiene.

Patients, HCWs and staff must cough or sneeze into their elbow sleeve or a tissue; throw away used tissues and immediately perform hand hygiene; refrain from touching their eyes, nose or mouth with unwashed hands. They must not share any food, drinks, unwashed utensils, cigarettes or vaping devices.

Point-of-care risk assessment (PCRA)

Prior to any patient interaction, all HCWs must assess the infectious risks posed to themselves, other HCWs, patients and visitors, the situation, or the procedure.

The PCRA is based on professional judgment about the clinical situation, as well as up-to-date information on how the specific health-care facility has designed and implemented appropriate physical (engineering) and administrative controls, and the use and availability of PPE.

To conduct a PCRA, evaluate the likelihood of exposure to COVID-19:

- **From a specific interaction** (e.g., performing or assisting with aerosol generating medical procedures (AGMPs), non-clinical interactions, direct face-to-face interactions with patients);
- **With a specific patient** (e.g., infants or young children, patients not able to practice hand hygiene or respiratory etiquette, frequent coughing or sneezing);
- **In a specific environment** (e.g., single rooms, hallways, assessment areas, etc. where physical distancing cannot be maintained).

See BCCDC for a sample [COVID-19 patient screening tool for health-care interactions & routine PCRA tool](#).

Guidance for Personal Protective Equipment

12. Key PPE recommendations

HCWs and non-clinical staff must follow provincial guidance for [Mask Use in Health-Care Facilities During the COVID-19 Pandemic](#) in all health-care settings.

Use eye protection for all HCWs who enter the patient/exam room or bed space or when within two metres of a patient.

For direct care of patients with suspected COVID-19 illness (e.g., patients with risk factors and/or symptoms of COVID-19) or a diagnosis of COVID-19, follow droplet and contact precautions. This includes wearing a medical mask, eye protection, gloves and gown.

Extend use of PPE wherever possible. This includes:

- In the same patient cohort¹, keep PPE on between patient encounters unless damaged or visibly soiled. The exception is gloves, which must be changed between each patient.
- Change PPE if moving from patients with confirmed COVID-19 to patients with suspected COVID-19.
- Change PPE if moving between patients on additional precautions for non-COVID-19 reasons (e.g., airborne, droplet and contact).
- Properly doff and dispose/clean and disinfect PPE when leaving the patient care area (e.g., at end of shift, during breaks or mealtimes).

Access to additional PPE, such as respirators, will be provided in circumstances where a HCW determines there is elevated risk of COVID-19 transmission through patient interaction.

When wearing PPE

- Avoid touching your mask or eye protection. If you must touch or adjust your mask or eye protection, perform hand hygiene immediately before and after adjusting.
- Leave the patient care area if you need to remove your mask (e.g., at end of shift or during a break) and ensure two metres distance from patient or exit exam room; do not re-use a doffed mask.
- Change PPE if it becomes damaged or visibly soiled.
- Use extreme care when taking off/doffing PPE and always perform hand hygiene after removing each individual piece of PPE and before putting on/donning new PPE.
- In the context of COVID-19, use of an N95 respirator or equivalent is **only** required when performing AGMPs on a person with suspected or confirmed COVID-19.
 - Use an N95 respirator or equivalent and eye protection (e.g., goggles or face shield), gloves and gown for AGMPs for patients with suspected or confirmed COVID-19.
 - In addition, adhere to routine institutional IPC and workplace safety guidelines and practices.
- Eye protection must be a well-fitting device that covers the front and sides of the face. Regular eyeglasses are **not** sufficient to protect from all splashes or droplet spray and are not considered adequate protection.
 - Eye protection, such as goggles, safety glasses or combination medical mask with attached visor, need to cover from the eyebrow to the cheekbone, and across from the nose to the boney area on the outside of the face and eyes. Eye protection should be fitted so that gaps between the edges of the eye protection and the face are kept to a minimum.

¹ **Patient cohort** refers to a group of patients with the same diagnosis or suspected diagnosis. In the case of COVID-19, patients with a confirmed COVID-19 diagnosis, patients suspected to have COVID-19 (diagnosis not yet confirmed), and patients without symptoms suggestive of COVID-19, can each be a respective patient cohort. Decisions regarding cohorting should be made in consultation between facility director/administrator, medical health officer or designate and client care leader.

- Full face shields should extend below the chin to cover the face, to the ears at both sides of the head, and there should be no exposed gap between the forehead and the shield's headpiece.
- For tasks with significant risk of splash, like AGMPs, a full-face shield or goggles must be used.
- When reusable eye protection is used for multiple patient encounters, it should be cleaned and disinfected as per the guidance found on [BCCDC's webpage](#).
- Properly doff, clean, and disinfect your eye protection when visibly soiled and when leaving the patient care area (e.g., at end of shift or during a break).

Putting on (donning) and taking off (doffing) PPE

For up-to-date information on PPE including putting on and taking off, as well as posters and signage, please refer to the [BCCDC's Personal Protective Equipment webpage](#).

Discontinuation of Droplet and Contact Precautions

For patients in the community, discontinuation of droplet and contact precautions should be aligned with the requirements for self-isolation as per [public health guidance for management of cases and contacts associated with COVID-19 in the community](#). This guidance supports a preference for a non-test-based strategy.

For management of patients in acute care settings, the strategies for discontinuation of additional precautions for patients may be different due to a risk assessment of their patient population. For these recommendations follow provincial [guidance for discontinuing additional precautions related to COVID-19 for admitted patients in acute care](#) and [decision tree](#).

13. PPE Guidance for Patients and Essential Visitors

See [mask use in health-care facilities during the COVID-19 pandemic](#) for guidance on mask use for patients and visitors in all health-care settings.

Essential visitors accompanying patients with symptoms of or a diagnosis of COVID-19 should follow droplet and contact precautions.

Patients and essential visitors must also follow routine practices and any additional precautions that are in place for non-COVID-19 reasons.

Office Management

14. Clinic Response Planning and Organization

It is useful to have clearly defined roles and responsibilities, balanced by cross-training of staff and planning for backfilling positions if a staff member is unable to work.

Develop or update a clinic response strategy to ensure that all staff roles are clearly defined, and that information and decision-making pathways are identified. Designate one qualified office member as the lead for coordinating COVID-19 response at the practice level, including staff responsibilities, information gathering and dissemination, and developing a preparedness plan for the clinic.

Please use the COVID-19 community clinic care preparedness checklist in [appendix A](#) of this document to aid planning.

General practitioners, nurse practitioners and physician specialists in community clinics can order PPE and critical supplies, including ABHR and disinfecting products through the Health PPE Portal.

Midwives continue to have access to PPE and critical supplies through the Product Distribution Centre website at the Ministry of Citizens' Services.

15. Staff Education and Communication

Develop or update a communication strategy to ensure HCWs and staff have the most up-to-date information. See the [key resources](#) section for links to the latest information on COVID-19.

Ensure staff have clear, up-to-date information for communicating with patients. Ensure there is a process for reporting health and safety concerns.

16. Psychosocial Support

BCCDC offers guidance for psychosocial planning for HCWs during the COVID-19 pandemic. Please see the [key resources](#) section of this document for further information.

17. Staff Scheduling and Reassignments

Staff scheduling should be considered when developing a coordinated pandemic response for the clinic:

- Consider adjusting clinic hours to accommodate patient and staffing needs, while supporting physical distancing and IPC measures.
- Assess employee availability when greater staffing needs, and employee absences for family or self-care are expected.

18. Sick Leave Policy

Have open and frank dialogue with all HCWs and staff about sick leave policy prior to any staff illness or time away from work due to self-isolation or quarantine.

Clearly communicate that HCWs and staff who have suspected or confirmed COVID-19 must self-isolate at home, as well as those who have had a test and are waiting for their results.

Some staff might be only mildly ill or already recovering and/or caring for others and still able to perform some of their duties remotely by Internet or telephone, depending on how a clinic is set up.

Key Resources

Information is available on the following topics relating to COVID-19:

- **Symptoms** of COVID-19: <http://www.bccdc.ca/health-info/diseases-conditions/covid-19/about-covid-19/symptoms>
- **Testing** for COVID-19: <http://www.bccdc.ca/health-professionals/clinical-resources/covid-19-care/covid-19-testing/viral-testing>
- **BC COVID-19 Self-Assessment Tool** can help determine the need for further assessment: <https://bc.thrive.health/>
- **Non-medical information** about COVID-19 is available 7:30 a.m. to 8:00 p.m., seven days a week at the following toll-free number: 1-888-COVID19 (1-888-268-4319).
- **HealthLink BC** and **8-1-1** for health advice on COVID-19 (translation services are available): <https://www.healthlinkbc.ca/health-feature/coronavirus-covid-19>
- **Information on masks:** <http://www.bccdc.ca/health-info/diseases-conditions/covid-19/prevention-risks/masks>
- **Information on COVID-19 immunization:**
[NACI Recommendations on the use of COVID-19 vaccines](#)
[BCCDC Monitoring vaccine update, safety and effectiveness \(March 30, 2021\)](#)
[BCCDC Getting a vaccine](#)

Other resources include:

- BCCDC's COVID-19 care webpage for HCWs: <http://www.bccdc.ca/health-professionals/clinical-resources/covid-19-care>
- COVID-19 signage and posters: <http://www.bccdc.ca/health-professionals/clinical-resources/covid-19-care/signage-posters>
- The Province of British Columbia's response to COVID-19: <https://www2.gov.bc.ca/gov/content/safety/emergency-preparedness-response-recovery/covid-19-provincial-support>
- Office of the Provincial Health Officer, orders and notices: <https://www2.gov.bc.ca/gov/content/health/about-bc-s-health-care-system/office-of-the-provincial-health-officer/current-health-topics/covid-19-novel-coronavirus>

Infection Prevention and Control resources:

- Public Health Agency of Canada: [Routine practices and additional precautions for preventing the transmission of infection in healthcare settings](#)
- BCCDC's poster on [environmental cleaning and disinfectants for clinic settings](#)
- B.C. Ministry of Health: [Best practice guidelines for cleaning, disinfection and sterilization of critical and semi-critical medical devices in BC health authorities](#)
- BCCDC's webpage on [personal protective equipment](#) for HCWs
- Vancouver Coastal Health's [IPAC best practices guideline: Point-of-care risk assessment](#).
- WorkSafeBC's [COVID-19 frequently asked questions: general ventilation and air circulation](#).

For Patient Management:

- BCCDC's webpage for information on [clinical care](#)

- BCCDC's [Interim Guidance: Public health management of cases and contacts associated with novel coronavirus \(COVID-19\) in the community](#).
- [Guidance for Primary Care Management of Adult Outpatients with Suspected or Confirmed COVID-19](#)
- BCCDC's [pregnancy](#) webpage for recommendations for antepartum, intrapartum and postpartum care for patients during the COVID-19 pandemic
- BCCDC's [patient handouts](#) for patient information

Support for HCWs:

- BCCDC's webpage on [exposures and return to work for health-care workers](#): Includes risk assessment and management for health-care workers exposed to COVID-19 patients and information on return to work after exposure or illness.
- BCCDC's [health-care provider support](#) webpage
- BCCDC's resource on psychological support: [Supporting the psychosocial well-being of health-care providers during the novel coronavirus \(COVID-19\) pandemic](#)
- Doctors of BC's [Physician Health Program \(PHP\)](#), which offers confidential advocacy, support and referral assistance for individual physicians and physicians-in-training. PHP also provides wellness initiatives to promote the ongoing health of our physician community.

References

BC Centre for Disease Control. Common Questions [Internet]. [cited 2020 Oct 30]. Available from: <http://www.bccdc.ca/health-info/diseases-conditions/covid-19/common-questions>

Mondelli MU, Colaneri M, Seminari EM, Baldanti F, Bruno R. Low risk of SARS-CoV-2 transmission by fomites in real-life conditions. *Lancet Infect Dis* [Internet]. 2020 Sep 29 [cited 2020 Nov 4]. Available from: [https://www.thelancet.com/journals/laninf/article/PIIS1473-3099\(20\)30678-2/abstract](https://www.thelancet.com/journals/laninf/article/PIIS1473-3099(20)30678-2/abstract)

Public Health Agency of Canada, Centre for Communicable Diseases and Infection Control (Canada). Hand hygiene practices in healthcare settings [Internet]. Ottawa, Ont.: Centre for Communicable Diseases and Infection Control, Public Health Agency of Canada; 2013 [cited 2020 May 14]. Available from: <https://www.deslibris.ca/ID/235647>

BC Ministry of Health. Pandemic Preparedness - Province of British Columbia [Internet]. [cited 2020 Apr 10]. Available from: <https://www2.gov.bc.ca/gov/content/health/about-bc-s-health-care-system/office-of-the-provincial-health-officer/current-health-topics/pandemic-influenza?keyword=2012&keyword=pandemic&keyword=plan>

Provincial Infection Control Network of BC. British Columbia Best Practices for Environmental Cleaning for Prevention and Control of Infections in All Healthcare Settings and Programs [Internet]. PICNet Guidelines and Toolkits. [cited 2020 May 10]. Available from: <https://www.picnet.ca/guidelines/environmental-cleaning/>

Daly P. Pandemic influenza and physician offices. *BCMJ*. 2007 Jun;49(No. 5):263–9.

Appendix A: COVID-19 Community Clinic Care Preparedness Checklist

COVID-19 Infection Prevention and Control (IPC) Preparedness Checklist for Community-Based Physicians, Nursing Professionals and Midwives Clinic Settings

General IPC Measures and Clinic Management

- Acquaint yourself with current clinical information about the recognition, treatment and prevention of transmission of COVID-19.
- Educate all staff about COVID-19.
- Make plans to ensure your family will be looked after in a pandemic while you are at work
- Develop a contingency plan for staff illnesses and shortages.
- Assign a staff member to coordinate pandemic planning and monitor public health advisories.
- Maintain copies of pandemic educational materials and self-care guides for patients (provided by public health).
- Appropriate COVID-19 [posters and signage](#) should be placed at entrance doors, reception area and exam rooms (preferably in all of these places) (e.g., facility entrance, medical mask and physical distancing posters).
- Post signage and create voicemail message advising patients to check in by phone before coming to in-person appointments.
- Post hand hygiene and cough etiquette signs in the waiting area.
- Ensure alcohol-based hand sanitizer (with at least 70% alcohol) is available at multiple locations: office entrance, reception counter, waiting room, and by every exam room for use before entering and upon exit.
- When available, provide staff with small bottles of alcohol-based hand sanitizer (with at least 70% alcohol).
- Consider installing physical barriers (e.g., plexiglass partitions) to separate patients from reception staff
- Limit use of shared items by patients (e.g., pens, clipboards, phones).
- Rearrange waiting room to ensure two-metre distance between people wherever possible.
- Remove difficult to clean items (e.g., toys) from the waiting area
- Replace cloth-covered furnishings with easy-to-clean furniture where possible.
- Provide disposable tissues and no-touch waste receptacles in waiting area and exam rooms.
- Provide plain soap and paper towels in patient washrooms and at staff sinks with clear instructions on hand hygiene.
- Display personal protective equipment (PPE) [putting on and taking off](#) instructions in locations available to all health-care staff.
- Empty exam rooms of all but bare minimum of equipment (e.g., exam table, chair, blood pressure cuff, lights).
- Provide paper sheeting for exam tables and change between patients.
- Keep frequently used doors open to avoid recurrent door handle contamination.

Patient and Staff Management

- Triage all patient requests for visits over the phone.
- In group practices, consider having one care provider or one team see all patients with suspect or confirmed COVID-19.
- If possible, schedule patients with symptoms suggestive of COVID-19 during designated time slots.
- If possible, provide a separate entrance and waiting area for patients with [symptoms](#) of COVID-19.
- Avoid multiple patients in the office at the same time and minimize the number of patients in waiting or exam rooms (e.g., patients to wait outside of the office or in their car until called in one at a time).
- Avoid non-essential visitors accompanying patients, where possible.
- Advise patients and accompanying essential visitors to practice diligent hand hygiene and respiratory etiquette.
- Provide patients and essential visitors with medical masks upon arrival, if available and medically tolerated.
- If possible, designate one exam room for all patients with symptoms suggestive of COVID-19, as close to the entrance as possible to minimize patient travel.
- Minimize the number of tasks that have to be done in the exam room (e.g., chart completion).
- Perform hand hygiene before and after each patient contact.
- Prior to any patient interaction, perform a PCRA to assess the likelihood of exposure to infectious agents.
- Medical mask use implemented for HCWs, non-clinical staff, patients, and visitors entering the facility in accordance with the [B.C. Ministry of Health policy communiqué: mask use in health-care facilities during the COVID-19 pandemic](#)
- Eye protection is used for all HCWs who enter the patient room or bed space or when within two metres of a patient.
- Wear recommended PPE for droplet and contact precautions (medical mask, eye protection, gown and gloves) for any direct contact with patients with suspected or confirmed COVID-19.
- Use an N95 respirator or equivalent and eye protection (e.g., goggles or face shield), gloves and gown for aerosol generating medical procedures (AGMPs) for patients with suspected or confirmed COVID-19.
- Properly doff and dispose of PPE when leaving patient care area (e.g., at end of shift or during a break) or when PPE is visibly soiled or damaged.
- Plan for disposition of all patients following office visit (choices will include):
 - Arrange testing per current guidelines (if suspected COVID-19)
 - Send home with self-care guide
 - Referral to alternate-care site
 - Admission to acute care
 - When referring patients with suspected or confirmed COVID-19, notify receiving facility in advance.
- Monitor staff illness and ensure staff with COVID-19 infection follow appropriate guidance. suspected or confirmed COVID-19.

Cleaning and Disinfection

- Inform all staff regarding current [cleaning and disinfection guidelines](#), including approved cleaning products.

- Clean and disinfect shared reusable medical equipment (e.g., stethoscopes, blood pressure cuffs, etc.) in between patients and at the end of each shift.
- Clean and disinfect exam rooms at least once a day (e.g., chairs, tables, floors).
- Clean and disinfect frequently touched surfaces at least twice a day (e.g., workstations, cell phones, doorknobs, etc.).
- Maintain a minimum two-week supply of plain soap, paper towels, hand sanitizer, cleaning supplies, and medical masks, if possible.

Note: This checklist is adapted from Daly, P. (2007). Pandemic influenza and physician offices [Electronic Version]