

COVID-19 Janssen Vaccine

Redistribution Guidelines

A. Introduction

This document provides information to support the internal redistribution of vaccine and will be updated to reflect best practices identified by health authorities and new information provided by vaccine manufacturers.

B. Key Considerations

1. Risk and responsibility for redistribution held by province, and health authorities.
2. Transport should only with unopened vials in a 2°C to 8°C environment; transport of pre-filled syringes may be done in exceptional circumstances (see section C.2).
3. An appropriate container must be used to transport vaccine.
4. Temperature must be monitored during transportation and documented.
5. Repacking should be done in a 2°C to 8°C environment whenever possible.
6. Transfers should be limited to decrease likelihood of temperature excursions and disruption to physical stability of vaccine.
7. Transfers between containers should be completed as quickly as possible.
8. Vials should be packaged as to minimize jostling.
9. Vaccine should not come into contact with ice packs.
10. Vaccine should be protected from light.
11. **The Janssen vaccine cannot be frozen.**

C. Methods of Redistribution

1. Redistribution of unopened vials (2°C to 8°C):

The Janssen vaccine should be transported using containers and temperature monitoring devices typically used for the transport of vaccines which must be maintained at temperatures of 2°C to 8°C. Considerations listed above should be fully applied and the time limits related to the stability of the vaccine at this temperature must be adhered to (see Section D).

Information on the provincial standards for shipping and receiving refrigerated vaccine can be found in the [Communicable Disease Control Manual Chapter 2 : Immunization Appendix E - Management of Biologicals](#).

Cold Chain Resources for Community Providers

- [How to Store Vaccines in the Refrigerator](#)
- [Packing an Insulated Cooler](#)
- [Handle Vaccines with Care](#)
- [Equipment Malfunction or Power Failure](#)
- [TempTale® Directions](#)
- [Temperature Form](#)
- [Cold Chain Checklist](#)

2. Distribution of prefilled syringes (2°C to 8°C, or up to 25°C or room temperature)

In exceptional circumstances, it is acceptable to pre-draw vaccine. Prefilled syringes should be carefully transported and labelled to ensure that they are not used outside of their window of stability.

After the first dose has been withdrawn, the filled syringe can be held at 2°C to 8°C for up to 6 hours or at room temperature (maximally 25°C) for up to 3 hours, after the first puncturing of the vial. Discard if vaccine is not used within this time.

Prefilled syringes unused at the end of a clinic may not be returned to the refrigerator for use the next day and must be wasted. In an effort to avoid wastage, care should be taken to pre-draw a reasonable number of doses.

D. Cold Chain Maintenance

- **Unopened** vials can be kept in a 2°C to 8°C environment until their expiry date.
- **Open vials** are stable in a 2°C to 8°C environment for up to a maximum of six hours after the first puncture of the vial.
- **Open vials** are stable at room temperature room temperature (maximally 25°C) for up to three hours after the first puncture of the vial.

E. Procurement of Cold Chain Equipment

All cold-chain equipment coordination and distribution will be handled centrally. This will include all refrigerated units, shipper containers, and data loggers. Equipment coordination questions and procurement requests should be directed to IBCOC_Operations@phsa.ca.

F. Selection of Data Loggers

The [Vaccine Storage and Handling Toolkit](#) (US CDC) has the following information regarding the selection of data loggers (they refer to them as DDL for “digital data logger”). Should health authorities wish to explore their own procurement of data loggers, any models considered should have the following specifications:

- Detachable probe that best reflects vaccine temperatures (e.g., a probe buffered with glycol, glass beads, sand, or Teflon®)
- Alarm for out-of-range temperatures
- Low-battery indicator
- Current, minimum, and maximum temperature display
- Recommended accuracy of +/-0.5°C (+/-1°F) or better
- Logging interval (or reading/recording rate) that can be programmed by the user to measure and record temperatures at least every 30 minutes
- Current and valid *Certificate of Calibration Testing*

G. Reporting Requirements

British Columbia is [required to report](#) back to the National Operations Centre information on the acceptability of all vaccine, and other materials received in province provided via a federal contract with the supplier. The provision of this information is critical for the federal government to monitor the implementation of the contracts they have established and supports the uninterrupted provision of vaccine and supplies in the future. Internal, provincial reporting requirements also require the provision of the following information:

Vaccine

1. Vaccine delivery site address
2. Purchase Order Number (if known)
3. Quantity received
4. Date of receipt
5. Time of Receipt
6. Damage to the packages? If yes, describe how many damaged and to what extent.
7. All Janssen boxes were received with no damage or missing vials (50 vials per single box).
8. Damage? If yes, describe how many and to what extent
9. Missing vials? Yes, how many?

Reporting Approach:

All sites receiving vaccine directly from the manufacturer should send a copy of their waybill and confirmation that the package was in tact directly to IBCOC_Operations@gov.bc.ca. If no waybill is included with the shipment, this information can be provided via email. This satisfies **data elements 1-6 above**.

Sites where trays are initially opened, allowing for the condition of the vials to be assessed, must provide **data elements 7-9 above** via email to IBCOC_Operations@phsa.ca.

H. Inventory Management

Regular inventory management processes should be followed. If questions or issues arise, consult with regional inventory management teams for advice and guidance.

Any questions or requests for revision
of this document should be sent to
IBCOC_Operations@phsa.ca

List of Amendments

Date	Section	Description	Author